1

Explor. in Capt Raynolds Dept [illegible] late

Nuphar polysepalum [illegible] foliis ovatis-oblongis

sinu angusto profunde cordatis; floris magni sub globosi sepalis 9–12 oviaris et medius maximis; petalis 12–18 spatulatis obtusis, staminibus numerosis; antheris apice truncato-appendiculatis; stamina demum recurvata aequantibus s. eis brevioribus; ovario urceolato striato stigmate peltato, radiis stigmatosis 13–21 disci umbilicati margine crenatum fere attingentibus, facca versus apicem constrictum nec rostratum subito. —

Hab. Lacus alpinas 6000–9000 ped alt. Montis Stopulorum

131

Inhabits small mountain lakes in the Rocky Mountains from Colorado to the sources of the Columbia River near Long's Peak, to those of the Platte 40 Lat 44

Dr F. V. Hayden collected it in the Yellowstone Lake Capt W. F. Raynolds Exp. June 20th 1860 in a small Lake between Henry's Fork and Snake Fork of the Columbia River at an altitude of 6500 ft

Miss Merrill brought from the Yellowstone 1862, a vague account of the plant and some specimens, to verify her tale, from Lake Gibson near Longs Peak, and finally Dr Parry gathered ample material in Osborne Lake 8800 ft above the sea near Longs Peak.

where it grows with Menyanthes trifoliata Utricularia intermedia, Scirpus, Carex etc. he found it in flower in August to the depth of the water being 5 ft. at the time

The Leaves are very much like those of N. luteum



Shipped, By Enno Sander

IN GOOD ORDER AND CONDITION, On account and risk of whom it may concern, on board the good **Steamboat** called the Key Stone Packet whereof H Johnson **is Master** for the present voyage, now lying at the Port of **Saint Louis,** and **bound** for ~~Key Stone~~ Ft Leavenworth the following packages or articles, marked or numbered as below, which are **to be delivered,** without delay, in like good order, at the said Port—Unavoidable dangers of the River and Fire only excepted—**unto** ~ or assigns, he or they **paying freight** for said goods at **the rate of** Freight Paid $2 per Sack and **Charges**

In Witness Whereof, The Owner, Master, or Clerk of said Steamboat, hath affirmed to three Bills of Lading, all of this tenor and date, one of which being accomplished, the others to stand void.

Dated at St. Louis, this 29th day of July 1864

Published and for sale by C. Witter, 27, South-east corner of Walnut and Second streets, St. Louis.

MARKS.	ARTICLES.	WEIGHT.
Jas P. Roy. Capt. & A. C.S. Ft Leavenworth For E. Jones. M. D. Care Medical Director Santa Fe N. M. Via St. L. & Ft. S. R.R.	1 Box Shot No 10. J. Campbell	52 lbs.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10 cm
copyright reserved
MO

2

than those of our N. advena, being more oval
in outline and ~~triangular~~ deltoid than orbicular, and with a narrower
more closed sinus, the obtuse lobes more gradually separating
from one another. In N. advena I find the sinus of
of 40 degrees and the lobes than acute triangular with
~~thrown~~ bent points. This form of the sinus and shape of the leaf
is by no means constant, for whenever the substance
of the leaf is more fully developed the lobes
become broader, more obtuse and the sinus of course
narrower, as I find it in specimens from Alleghany
while sometimes as in specimens from Houston
Texas, the sinus becomes closed up and the
lobes even overlap. our species were floating when observed.

Those leaves of N. sagittifolia Nuttall
8 1/2 - 9 1/2 inches long and 6 1/2 - 7 1/2 inches
wide, ~~have the~~ ~~an average~~ These 5 leaves gave
their exponent and ~~average~~ the same rule
of leaves of N. lutea give each an average
exponent of length and width = 10 : 8. -
Each the same number of leaves of N. advena
from different parts of the country gave the
exponent = 10 : 9. The difference is appreciable
small enough, but in the appearance of the
leaf quite striking. I find also a difference
in the venation of the leaves of these species

130

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

THE STATE OF MISSOURI, } ss.

COUNTY OF ST. LOUIS,

To Dr. Geo Engelmann Greeting:

You are hereby commanded that, setting aside all manner of excuse and delay, you appear before our **ST. LOUIS COURT OF COMMON PLEAS**, at the City of St. Louis, on the 12th day of June 1865, at 10 o'clock, a M., then and there to testify, and the truth to say, in a certain cause pending in said Court, wherein Wm. Ballentine plaintiff and North Missouri Rail Road Company defendant on the part of dept and herein you are in no wise to fail.

GEORGE ENGELMANN PAPERS
MISSOURI BOTANICAL GARDEN

Witness, LEON J. PAPIN, Clerk of our said Court, with the seal thereof hereto affixed, at office, in the City of St. Louis, this 10th day of June in the year of our Lord eighteen hundred and sixty-five

Leon J. Papin, Clerk.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

3 there being nearly 3 times as many veins ~~counted~~ with the midrib as there ~~were counted~~ at the base, while in both N. lutea and N. advena I noticed only about twice as many for the midrib as from the base. It may not be out of place to note in connection that the distribution of the veins of the species of Nymphaea which I have been able to ~~compare~~, can be readily distinguished from all those of Nymphaea so far as my observations extend; the former having by far the largest number of secondary ribs connected with the midrib, and ~~usually~~, while in Nymphaea most ribs are basilar, and few only come from the midrib. The 129 plant a little emersed

The flowers of our species are the largest of any known Nuphar, and are composed of 12 sepals, ~~the~~ many more petals in ~~any other species~~ whence the name,

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

St. Louis 17th March 1865

George Engelmann, M. D.

Dear Sir

You are respectfully requested to attend a regular meeting of the Board of Trustees of the St. Louis College of Pharmacy at the Room of the St. Louis Medical Faculty.

on Tuesday Evening March 21st at half past 7 o'clock P. M.

Punctual attendance is requested.

By order of the chairman

James McBride

Sec't.

4/ the sepals being concave ~~are~~ ~~partly~~ ~~rolled~~ ~~over~~ the ~~petals~~ "the inner ones curved in" "partially conceal the greater part of the purple inner petals"
~~corpse~~, the shape of the flower is ~~urceolate~~ globular (cupshaped)
somethg like 3 inches in diameter, but when fully
laid open it measures 4½ to 5 inches, "
~~diameter~~ The sepals are arranged not as it was
at first supposed in 3 or 4 whorls of 3 sepals each
but in ~~4~~ 5/13 arrangt, or perhaps, the outer ones
in 2/5 and gradually changing in to 5/13 ~~position~~
divergence; the sepals increase in size & petaloid
appearance from the outermost or first to the 7 or
8th when the decrease again in size, but become
of more delicate petaloid structure and colour
than the outermost ones are oval 1¼ - 1½ inches long, green
with yellowish margin; the next 2 or 3 are clear
obovate-oblong, 2 - 2½ inches long, and of a yellowish
colour green at base, and often with the peach blossom
the following ones are the largest & of a transverse "spoon-
form, obovate-oblong, yellow, with the peach blossom
and purple edge, obovate-spatulate 1½ - 1 inches long

especially on the outer edge

5 Yellow or Reddish Brown

Dr Parry informs me that he
recollects that this is
there are often. taken these seeds with the
petals. The petals themselves are spatulate truncate
12 to 18 in number, 9 l. long - 6 l. broad (Parry)
-th dry specimens of this the only 5 l. long - from 2-4
lines wide, gradually narrowed to the claw 127
they are deep yellow, or in Dr Parry's specimens, deep
red in the middle & yellow at base & tip, the claws
are more numerous than the middle ones, & together with the appendage are deep red
Relieved by the bright yellow (coral pink tips as in
the genus) Rollen the outer ones shorter & broader
the inner ones longer & narrower; at maturity they
spread or recurved
deeply umbilicate &
Margin dor. usually with 15-24
usually (usually 16-19) stigmatic rays with
extend to near the (regularly) margin. The yellowish
green or deep red or red (with yellow) edged
see note #
smooth & glossy, deep
Dr Parry considers only half ripe fruit, the ripe
& ripened specimens brought by
brought by us (dry) 1 3/4 long & 1/2 - diameter

ALEX. LEITCH.

APOTHECARY & PHARMACEUTIST,

Marble Building Cor. Fourth & Olive Sts. St. Louis, Mo.



134

0 cm 1 2 3 4 5 6 7 8 9 10
copyright reserved
MO

Nuphars
polysepalum
June 15th 1865

5/13 arrangt
of sepals
verte



MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

dist with the dried specimen before me 9–11 lines in diameter :—

(the only one that I dissected) — Unlike
the dist in the wetter specimen of N. adnata DC —
the dist entire and the rays of the stigma not
extended to meet the margin, while in N. lutea
they do reach nearly to the annulate cilia
just as in N. stricta polyscepha —

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

GEORGE EASTMAN HOUSE PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

Enno Sander,

Ecke der Zweiten und Walnut Straße,

unter Barnum's Hotel,

Saint Louis, Mo.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN HERBARIUM

0 cm 1 2 3 4 5 6 7 8 9 10
Copyright reserved
MO



0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

Enno Sander,

Ecke der Zweiten und Walnut Straße,

unter Barnum's Hotel,

Saint Louis, Mo.

MISSOURI BOTANICAL
GEORGE ENGELMANN

MO
cm 0 1 2 3 4 5 6 7 8 9 10
copyright reserved

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

Detroit

Nuphar advena

November 1865

from Park Marsh

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

(4 cm)

Wood Nelson 1850 ~~1850~~ 1900 along the road to Mt. Diablo

— across — 3-15 ~~then~~ then turning up hill to the
(3 miles) house at the end of the lane
(on hill overlooking valley)

— Nelson — 8 miles to the mouth of Diablo
(2 cm)

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

Emil Sander,

Ecke der Zweiten und Walnut Straße,

unter Barnum's Hotel,

Saint Louis, Mo.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN FUND

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

Nephrodium
July IX 64

142

Nephrodium dilatatum, sagittifolium with 6 spots
Cuba Kalmia
sagittifolia has no spots

Alex. Leitch,

APOTHECARY & CHEMIST,

Cor. 4th & Olive Sts., St. Louis.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

MO
0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved

Aspidium Kaduricum ? Bible 25th July 20th 1865
Mostly 6 lobed - one Specimen 8 lobed - colour
purple at the base internally - leaves glossy - very
thin - lobes approximate rounded

M. Variegatum

141

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

upper variegation

Section of petiole

Detail
sheet 7

MISSOURI BOTANICAL GARDEN
GEORGE E.

Rec'd Jan 21st

Copy

Notes on Nuphar pictum

Stigmatic rays irregular in number 13 – 19 of 12 Specimens examined 4 had 14 rays 4 19 rays 3 had 16 rays 1 had 13 rays the larger plants having the largest number; The rays do not extend to the margin an often variable in length and as a general rule point in the direction outward of the rounded points of the irregular crenated edge of the stigma; Entire ovary sulcate umbilicate perpendicularly ridged on the outside by ridges corresponding to the projecting points of the crenated edge and the stigmatic rays

Petals broad spatulate (slightly concave) 12 to 16 somewhat variable in width yellow at their attachment & the ovary deep red towards the center and yellow at the tips, about 3/4 inch long 1/2 inch broad projecting slightly beyond the 4th row of Stamens

Sepals in threes regularly in 3 circles alternating, rarely in 4 circles or with one or more petals irregularly developed –, outer smaller irregularly scoop shaped green with the margin inclined to yellow, next inner row of sepals larger broad orbicular - spatulate, with a wide attachment at base, more yellow than the outside sepals but the lower outside projection light green. (less concave than the outer row)

over

0 1 2 3 4 5 6 7 8 9 10 cm
copyright reserved
MO

The sections of the stems also should be preserved and attached to the respective flowers with the pressed stem projecting into the bottles.

The writer is a subscriber of your valuable book of observations, and in the spirit and from the love of the subject (which I have in common with many boys) and took notes, studied to press dried specimens and write down

I hope you will find it agreeable to have these observations and specimens to examine from time to time, and that you will be encouraged and prepared to receive and preserve all material given

Yours truly

C C Starr

Davenport Iowa 15th Feb 1865

Missouri Botanical Garden
George Engelmann Papers



; inner sepals generally narrower & shorter with a
redish tinge mostly on the outer edge. (which becomes
more conspicuous in fading) in color resembling
the deep blush of a rosey peach. (The whole flower
is globular cup shaped. the upper edge of the sepals
curving inward & concealing the greater part of
the dense mass of stamens.) (P.S. Something like this
Sort of the first named species will give you the
best idea. I suppose the dried specimen would be somewhat
shrunk but you can probably make due allowance. I
compare in coloring &c with dried specimens of
the former; Anthers linear spatulate nearly as long as
the filaments of a deep red color the upper surface
pollen-form & coated with the yellow powder of pollen
the anthers curving outward and forming a dense
deep red rosulate disc in full flower specimen
reaching up to the inner edge of the stigmatic disc
the developing ovary extending beyond and hiding the
withered remains of anthers petals & sepals closely
attached at the base. Fruit. (see specimen sent you)
furrowed especially towards the top of a deep green color
smooth & glossy.. Seeds (mature) enveloped in
shining cellular tissue (about a line in diameter)

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

Set 1 from Belle Isle Nov 1865

Right Valvula

138

upper side

under surface



0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

Detroit from Bloody Run Nov 1865

Heights various

139

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

MISSOURI BOTANICAL
GEORGE ENGELMANN

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

145

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

copyright reserved

81-4

Nymphaea

cm 0 1 2 3 4 5 6 7 8 9 10
copyright reserved
MO

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

Gray Writes Aug 28 1865

The downy Nymphaea from Central N York is

N. mirabor DC. syst.

Paine of Utica has just gathered a <u>seedless</u>
Nymphaea near Lake Ontario. leaves immensely
large, veins very prominent beneath; therefore
probably N. uniformis DC. flowers apparently
= odorata, but odorless or almost entirely so.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

Allenton, St. Louis Co. Mo.
Nobr. 14th, 1864.

Lieber Hr. Doctor!

Freilich

schicke ich Ihnen meine meteorologischen Beobachtungen für den Monat October 1864. Wir sind somit wohl gesund und mit Beobachtungen der Vögel und sonstigen Dingen beschäftigt. Sonst giebts nichts Neues hier. Heute schicke Ihnen wir doch Ihre Beobachtungen für Septbr. und October 1864, damit sie sie mit den meinigen vergleichen kann.

Ganz ergebenst.

H. H.

A. Fendler.

George Engelmann M.D.
St Louis, Mo.

Aug 25 1916

Antirrhinum hair on
petiole of Nymphaea
stem by Group from Ithica
N York

Hair ½ — 2 mm long.

Length of Pore — $\frac{10-14}{280}$

= 0.036 — 0.050 mm

X 60

Base ½
of hair X 280

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

ALEX. LEITCH.

APOTHECARY & PHARMACEUTIST,

Marble Building Cor. Fourth & Olive Sts. St. Louis, Mo.

cm 0 1 2 3 4 5 6 7 8 9 10
Copyright reserved
MO

Nymphaea elegans Hook coll. by Wright
on the Rio Grande, sent by Hooker
from Wright's seeds. -
Nymphaea amplas ? Wright Rio Grande, Dentate leaves

148

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

Nymphaea mikrosperma Rhizoma gracili ramoso profunde repente; foliis natantibus (ovatis subcordatis margine crenatis) minoribus cordatis orbiculatis seu ellipticis orbiculatis, (lobis obtusis, angulatis s. obtuse acutulatis, subtus rubellis seu nervis) parum prominulis, appendiculatis, sinu clauso sinu angusto, rarius clauso; stipulis latitudine longioribus ovatis margine membranaceis, pedunculis teretibus foliosis pilosulis versus petiolos (pedunculisque) villosis, floribus minoribus (albis s. extus pallide roseis) odoratissimis V. odoratissimi extensoribus obconicotubo elongato cuspidatis, interioribus antheris, pollinis stigmate . . . baccâ globosa, seminibus totis + arillum laxum involventibus.

But if only a variety of N. odorata I would say:

var. minor, Sims.: foliis subtus rubellis nervis minus prominentibus, sinu magis clauso, petiolis pedunculisque versus apicem villosis; floribus minoribus saepe extus roseis.

Letter to Vaux Nov 6 1865

0 1 2 3 4 5 6 7 8 9 10 cm copyright reserved MO

DEALER IN
PHARMACEUTICAL
Preparations,
FINE CHEMICALS,
And all the
New Remedies.

E. L. MASSOT,
CORNER OF
Fourth and Spruce Streets,
ST. LOUIS.

DISPENSING
AND
MANUFACTURING
PHARMACEUTIST.

ALER IN
ACEUTICAL
parations,
EMICALS,
d all the
Remedies.

E. L. MASSOT,
CORNER OF
Fourth and Spruce Streets,
ST. LOUIS.

DI
MAN
PHA

cm 0 1 2 3 4 5 6 7 8 9 10
copyright reserved
MO

Nymphaea alba

Casp. Schr. Ges. Naturf. Berl. 1855 p. 26

I. Melocarpa

A d'pressa

a chlorocarpa

N. venusta Hedge

N. rotundifolia Hedge

b. erythrocarpa

N. erythrocarpa Hedge

B circumvallata

C. sphaerocarpa

a chlorocarpa

N. carnea Hedge

b. splendens

N. splendens Hedge

b. erythrocarpa

D recurvata

N. recurvata Hedge

II. Oocarpa

N. candida Presl

N. candidissima Lehmann

N. semiaperta Klinggraeff

N. neglecta Hanslemann

N. Kosteletzkii Fallaux

N. intermedia Weber

N. pygmaea Hedge

N. quadrata Kurr et Kirk.

N. Gladstoniana Jacq.

F. aperta

F semiaperta

G. aperta

0 1 2 3 4 5 6 7 8 9 10 cm
copyright reserved
MO

M. W. ALEXANDER,

South-east corner Fourth and Market Streets,

ST. LOUIS.

---

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

Hypsiglena ochrorhyncha

Ouray, Uintah Co.

Nov 8 1955

$\frac{14}{405}$ 0.034 mm

151

$\frac{245}{745}$ = 0.031

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

...EX LEITCH

...IECARY & PHARMACEUTIST,

...R. 4TH & OLIVE STS., ST. LOUIS.

0 cm 1 2 3 4 5 6 7 8 9 10
copyright reserved
MO

Hyoptera tobaccoides Rau & Wiss.
Caiuá Desert, SP

$\frac{15}{415} = .036$ m

152

$\frac{28}{785} = .036$

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

...EX ...LEITCH

...ECARY & PHARMACEUTIST,

4TH & OLIVE STS., ST. LOUIS.

cm 0 1 2 3 4 5 6 7 8 9 10
copyright reserved
MO

Nymphaea odorata ?

St Louis 1838

Nov 8 1865

Pollen x 415



$\frac{17}{415} = 0.041$ mm

153

Pollen x 785

$\frac{315}{7850} = 0.040$

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

...EX. LEITCH

...ECARY & PHARMACEUTIST,

...4TH & OLIVE STS., ST. LOUIS.

cm 0 1 2 3 4 5 6 7 8 9 10
copyright reserved
MO

Nymphaea alba

Linkfeld 1825

Nov B 1865

Collen × 415



$\frac{16}{415} = 0.039$ mm

154

$\frac{316}{7850} = 0.040$

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MO

...ex Lerch

...HECARY & PHARMACEUTIST,

...R. 4TH & OLIVE STS., ST. LOUIS.

cm 0 1 2 3 4 5 6 7 8 9 10
copyright reserved
MO